

INFORMATION IN THE TABLE BELOW AND IN THE TABLES ON PAGES 3-5 MAY BE USEFUL IN ANSWERING THE QUESTIONS IN THIS SECTION OF THE EXAMINATION.

DO NOT DETACH FROM BOOK.

PERIODIC TABLE OF THE ELEMENTS

1	2																	3	4	5	6	7	8	9	10									
H 1.008	He 4.00																	B 10.81	C 12.01	N 14.01	O 16.00	F 19.00	Ne 20.18											
3																		13	14	15	16	17	18											
Li 6.94	Be 9.01																	Al	Si	P	S	Cl	Ar											
11	12																	26.98	28.09	30.97	32.06	35.45	39.95											
Na	Mg 24.30																	29	30	31	32	33	34	35	36									
19	20																	Ga	Zn	Ge	As	Se	Br	Kr										
K 39.10	Ca 40.08																	69.72	65.39	72.59	74.92	78.96	79.90	83.80										
37	38																	48	47	46	45	44	43	42	41	40								
Rb	Sr																	Cd	In	Sn	Sb	Te	I	Xe										
85.47	87.62																	112.41	114.82	118.71	121.75	127.60	126.91	131.29										
55	56																	80	81	82	83	84	85	86										
Cs	Ba																	Hg	Tl	Pb	Bi	Po	At	Rn										
132.91	137.33																	200.59	204.38	207.2	208.98	(209)	(210)	(222)										
87	88																	79	78	77	76	75	74	73	72	71								
Fr (223)	Ra 226.02																	Au	Pt	Ir	Os	Re	W	Ta	Hf	Rf								
																		196.97	195.08	192.2	190.2	186.21	183.85	180.95	178.49	178.49								
																		272	(271)	(268)	(277)	(264)	(266)	(262)	(261)	(261)								
																		Rg	Ds	Mt	Hs	Bh	Sg	Db	Rf	Ac								
																		(272)	(271)	(268)	(277)	(264)	(266)	(262)	(261)	227.03								
																		111	110	109	108	107	106	105	104	89								
																		Lanthanide Series																
																		196.97	195.08	192.2	190.2	186.21	183.85	180.95	178.49	178.49	138.91							
																		200.59	204.38	207.2	208.98	(209)	(210)	(222)										
																		251	252	257	258	259	262											
																		Cf	Es	Fm	Md	No	Lr											
																		(251)	(252)	(257)	(258)	(259)	(262)											
																		97	96	95	94	93	92	91	90	89								
																		Bk	Cm	Am	Pu	Np	U	Pa	Th	Ac								
																		(247)	(247)	(243)	(244)	(237)	238.03	231.04	232.04	227.03								
																		111	110	109	108	107	106	105	104	89								
																		Lanthanide Series																
																		196.97	195.08	192.2	190.2	186.21	183.85	180.95	178.49	178.49	138.91							
																		200.59	204.38	207.2	208.98	(209)	(210)	(222)										
																		251	252	257	258	259	262											
																		Cf	Es	Fm	Md	No	Lr											
																		(251)	(252)	(257)	(258)	(259)	(262)											
																		97	96	95	94	93	92	91	90	89								
																		Bk	Cm	Am	Pu	Np	U	Pa	Th	Ac								
																		(247)	(247)	(243)	(244)	(237)	238.03	231.04	232.04	227.03								
																		111	110	109	108	107	106	105	104	89								
																		Lanthanide Series																
																		196.97	195.08	192.2	190.2	186.21	183.85	180.95	178.49	178.49	138.91							
																		200.59	204.38	207.2	208.98	(209)	(210)	(222)										
																		251	252	257	258	259	262											
																		Cf	Es	Fm	Md	No	Lr											
																		(251)	(252)	(257)	(258)	(259)	(262)											

*Lanthanide Series

† Actinide Series

STANDARD REDUCTION POTENTIALS IN AQUEOUS SOLUTION AT 25°C

Half-reaction		$E^\circ(\text{V})$
$\text{F}_2(\text{g}) + 2\text{e}^-$	$\rightarrow 2\text{F}^-$	2.87
$\text{Co}^{3+} + \text{e}^-$	$\rightarrow \text{Co}^{2+}$	1.82
$\text{Au}^{3+} + 3\text{e}^-$	$\rightarrow \text{Au}(\text{s})$	1.50
$\text{Cl}_2(\text{g}) + 2\text{e}^-$	$\rightarrow 2\text{Cl}^-$	1.36
$\text{O}_2(\text{g}) + 4\text{H}^+ + 4\text{e}^-$	$\rightarrow 2\text{H}_2\text{O}(\text{l})$	1.23
$\text{Br}_2(\text{l}) + 2\text{e}^-$	$\rightarrow 2\text{Br}^-$	1.07
$2\text{Hg}^{2+} + 2\text{e}^-$	$\rightarrow \text{Hg}_2^{2+}$	0.92
$\text{Hg}^{2+} + 2\text{e}^-$	$\rightarrow \text{Hg}(\text{l})$	0.85
$\text{Ag}^+ + \text{e}^-$	$\rightarrow \text{Ag}(\text{s})$	0.80
$\text{Hg}_2^{2+} + 2\text{e}^-$	$\rightarrow 2\text{Hg}(\text{l})$	0.79
$\text{Fe}^{3+} + \text{e}^-$	$\rightarrow \text{Fe}^{2+}$	0.77
$\text{I}_2(\text{s}) + 2\text{e}^-$	$\rightarrow 2\text{I}^-$	0.53
$\text{Cu}^+ + \text{e}^-$	$\rightarrow \text{Cu}(\text{s})$	0.52
$\text{Cu}^{2+} + 2\text{e}^-$	$\rightarrow \text{Cu}(\text{s})$	0.34
$\text{Cu}^{2+} + \text{e}^-$	$\rightarrow \text{Cu}^+$	0.15
$\text{Sn}^{4+} + 2\text{e}^-$	$\rightarrow \text{Sn}^{2+}$	0.15
$\text{S}(\text{s}) + 2\text{H}^+ + 2\text{e}^-$	$\rightarrow \text{H}_2\text{S}(\text{g})$	0.14
$2\text{H}^+ + 2\text{e}^-$	$\rightarrow \text{H}_2(\text{g})$	0.00
$\text{Pb}^{2+} + 2\text{e}^-$	$\rightarrow \text{Pb}(\text{s})$	-0.13
$\text{Sn}^{2+} + 2\text{e}^-$	$\rightarrow \text{Sn}(\text{s})$	-0.14
$\text{Ni}^{2+} + 2\text{e}^-$	$\rightarrow \text{Ni}(\text{s})$	-0.25
$\text{Co}^{2+} + 2\text{e}^-$	$\rightarrow \text{Co}(\text{s})$	-0.28
$\text{Cd}^{2+} + 2\text{e}^-$	$\rightarrow \text{Cd}(\text{s})$	-0.40
$\text{Cr}^{3+} + \text{e}^-$	$\rightarrow \text{Cr}^{2+}$	-0.41
$\text{Fe}^{2+} + 2\text{e}^-$	$\rightarrow \text{Fe}(\text{s})$	-0.44
$\text{Cr}^{3+} + 3\text{e}^-$	$\rightarrow \text{Cr}(\text{s})$	-0.74
$\text{Zn}^{2+} + 2\text{e}^-$	$\rightarrow \text{Zn}(\text{s})$	-0.76
$2\text{H}_2\text{O}(\text{l}) + 2\text{e}^-$	$\rightarrow \text{H}_2(\text{g}) + 2\text{OH}^-$	-0.83
$\text{Mn}^{2+} + 2\text{e}^-$	$\rightarrow \text{Mn}(\text{s})$	-1.18
$\text{Al}^{3+} + 3\text{e}^-$	$\rightarrow \text{Al}(\text{s})$	-1.66
$\text{Be}^{2+} + 2\text{e}^-$	$\rightarrow \text{Be}(\text{s})$	-1.70
$\text{Mg}^{2+} + 2\text{e}^-$	$\rightarrow \text{Mg}(\text{s})$	-2.37
$\text{Na}^+ + \text{e}^-$	$\rightarrow \text{Na}(\text{s})$	-2.71
$\text{Ca}^{2+} + 2\text{e}^-$	$\rightarrow \text{Ca}(\text{s})$	-2.87
$\text{Sr}^{2+} + 2\text{e}^-$	$\rightarrow \text{Sr}(\text{s})$	-2.89
$\text{Ba}^{2+} + 2\text{e}^-$	$\rightarrow \text{Ba}(\text{s})$	-2.90
$\text{Rb}^+ + \text{e}^-$	$\rightarrow \text{Rb}(\text{s})$	-2.92
$\text{K}^+ + \text{e}^-$	$\rightarrow \text{K}(\text{s})$	-2.92
$\text{Cs}^+ + \text{e}^-$	$\rightarrow \text{Cs}(\text{s})$	-2.92
$\text{Li}^+ + \text{e}^-$	$\rightarrow \text{Li}(\text{s})$	-3.05

ADVANCED PLACEMENT CHEMISTRY EQUATIONS AND CONSTANTS

ATOMIC STRUCTURE

$$E = h\nu \quad c = \lambda\nu$$

$$\lambda = \frac{h}{m\nu} \quad p = m\nu$$

$$E_n = \frac{-2.178 \times 10^{-18}}{n^2} \text{ joule}$$

EQUILIBRIUM

$$K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$$

$$K_b = \frac{[\text{OH}^-][\text{HB}^+]}{[\text{B}]}$$

$$K_w = [\text{OH}^-][\text{H}^+] = 1.0 \times 10^{-14} \text{ @ } 25^\circ\text{C}$$

$$= K_a \times K_b$$

$$\text{pH} = -\log[\text{H}^+], \text{pOH} = -\log[\text{OH}^-]$$

$$14 = \text{pH} + \text{pOH}$$

$$\text{pH} = \text{p}K_a + \log \frac{[\text{A}^-]}{[\text{HA}]}$$

$$\text{pOH} = \text{p}K_b + \log \frac{[\text{HB}^+]}{[\text{B}]}$$

$$\text{p}K_a = -\log K_a, \text{p}K_b = -\log K_b$$

$$K_p = K_c(RT)^{\Delta n},$$

where Δn = moles product gas – moles reactant gas

THERMOCHEMISTRY/KINETICS

$$\Delta S^\circ = \sum S^\circ \text{ products} - \sum S^\circ \text{ reactants}$$

$$\Delta H^\circ = \sum \Delta H_f^\circ \text{ products} - \sum \Delta H_f^\circ \text{ reactants}$$

$$\Delta G^\circ = \sum \Delta G_f^\circ \text{ products} - \sum \Delta G_f^\circ \text{ reactants}$$

$$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$$

$$= -RT \ln K = -2.303 RT \log K$$

$$= -n\mathcal{F}E^\circ$$

$$\Delta G = \Delta G^\circ + RT \ln Q = \Delta G^\circ + 2.303 RT \log Q$$

$$q = mc\Delta T$$

$$C_p = \frac{\Delta H}{\Delta T}$$

$$\ln[A]_t - \ln[A]_0 = -kt$$

$$\frac{1}{[A]_t} - \frac{1}{[A]_0} = kt$$

$$\ln k = \frac{-E_a}{R} \left(\frac{1}{T} \right) + \ln A$$

E = energy v = velocity
 ν = frequency n = principal quantum number
 λ = wavelength m = mass
 p = momentum

Speed of light, $c = 3.0 \times 10^8 \text{ m s}^{-1}$

Planck's constant, $h = 6.63 \times 10^{-34} \text{ J s}$

Boltzmann's constant, $k = 1.38 \times 10^{-23} \text{ J K}^{-1}$

Avogadro's number = $6.022 \times 10^{23} \text{ mol}^{-1}$

Electron charge, $e = -1.602 \times 10^{-19} \text{ coulomb}$

1 electron volt per atom = 96.5 kJ mol^{-1}

Equilibrium Constants

K_a (weak acid)
 K_b (weak base)
 K_w (water)
 K_p (gas pressure)
 K_c (molar concentrations)

S° = standard entropy
 H° = standard enthalpy
 G° = standard free energy
 E° = standard reduction potential
 T = temperature
 n = moles
 m = mass
 q = heat
 c = specific heat capacity
 C_p = molar heat capacity at constant pressure
 E_a = activation energy
 k = rate constant
 A = frequency factor

Faraday's constant, $\mathcal{F} = 96,500 \text{ coulombs per mole of electrons}$

Gas constant, $R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$
 $= 0.0821 \text{ L atm mol}^{-1} \text{ K}^{-1}$
 $= 62.4 \text{ L torr mol}^{-1} \text{ K}^{-1}$
 $= 8.31 \text{ volt coulomb mol}^{-1} \text{ K}^{-1}$

GASES, LIQUIDS, AND SOLUTIONS

$$PV = nRT$$

$$\left(P + \frac{n^2a}{V^2}\right)(V - nb) = nRT$$

$$P_A = P_{total} \times X_A, \text{ where } X_A = \frac{\text{moles A}}{\text{total moles}}$$

$$P_{total} = P_A + P_B + P_C + \dots$$

$$n = \frac{m}{M}$$

$$K = ^\circ\text{C} + 273$$

$$\frac{P_1V_1}{T_1} = \frac{P_2V_2}{T_2}$$

$$D = \frac{m}{V}$$

$$u_{rms} = \sqrt{\frac{3kT}{m}} = \sqrt{\frac{3RT}{M}}$$

$$KE \text{ per molecule} = \frac{1}{2}mv^2$$

$$KE \text{ per mole} = \frac{3}{2}RT$$

$$\frac{r_1}{r_2} = \sqrt{\frac{M_2}{M_1}}$$

molarity, M = moles solute per liter solution

molality = moles solute per kilogram solvent

$$\Delta T_f = iK_f \times \text{molality}$$

$$\Delta T_b = iK_b \times \text{molality}$$

$$\pi = iMRT$$

$$A = abc$$

P = pressure

V = volume

T = temperature

n = number of moles

D = density

m = mass

v = velocity

u_{rms} = root-mean-square speed

KE = kinetic energy

r = rate of effusion

M = molar mass

π = osmotic pressure

i = van't Hoff factor

K_f = molal freezing-point depression constant

K_b = molal boiling-point elevation constant

A = absorbance

a = molar absorptivity

b = path length

c = concentration

Q = reaction quotient

I = current (amperes)

q = charge (coulombs)

t = time (seconds)

E° = standard reduction potential

K = equilibrium constant

OXIDATION-REDUCTION; ELECTROCHEMISTRY

$$Q = \frac{[C]^c [D]^d}{[A]^a [B]^b}, \text{ where } aA + bB \rightarrow cC + dD$$

$$I = \frac{q}{t}$$

$$E_{\text{cell}} = E_{\text{cell}}^\circ - \frac{RT}{n\mathcal{F}} \ln Q = E_{\text{cell}}^\circ - \frac{0.0592}{n} \log Q \text{ @ } 25^\circ\text{C}$$

$$\log K = \frac{nE^\circ}{0.0592}$$

$$\text{Gas constant, } R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$$

$$= 0.0821 \text{ L atm mol}^{-1} \text{ K}^{-1}$$

$$= 62.4 \text{ L torr mol}^{-1} \text{ K}^{-1}$$

$$= 8.31 \text{ volt coulomb mol}^{-1} \text{ K}^{-1}$$

$$\text{Boltzmann's constant, } k = 1.38 \times 10^{-23} \text{ J K}^{-1}$$

$$K_f \text{ for H}_2\text{O} = 1.86 \text{ K kg mol}^{-1}$$

$$K_b \text{ for H}_2\text{O} = 0.512 \text{ K kg mol}^{-1}$$

$$1 \text{ atm} = 760 \text{ mm Hg}$$

$$= 760 \text{ torr}$$

$$\text{STP} = 0.00^\circ\text{C and } 1.0 \text{ atm}$$

$$\text{Faraday's constant, } \mathcal{F} = 96,500 \text{ coulombs per mole of electrons}$$